Applicants: Philip O. Livingston and Friedhelm Helling

Serial No.: 08/475,784 Filed: June 7, 1995

Page 2

## Amendments to the Brief Description of the Figures

Please replace the paragraph beginning at page 5, line 4, with the following amended paragraph:

Figure 1A-1B

1A and 1B: The synthesis of GD3
protein conjugates after ozone
cleavage and reductive amination.
Insert represents HPTLC of GD3
before (lane A) and after (lane B)

the cleavage.

Please replace the paragraph beginning at page 6, line 11, with the following amended paragraph:

Figures 8A 1 and 8A 2 8A - 8B

8A and 8B: Specificity of peak titer sera from patients immunized GM2-KLH + QS-21 vaccine determined by immune chromatography layer described previously (3, Reference of the Third Series of Experiments). GM2 (A) and melanoma tissue ganglioside extract (B) were applied to TLC plates, incubated with sera from individual patients and stained with peroxidase-labeled goat antihuman IgM or IgG antibody. MAb 696 was used as positive control for GM2 and resorcinol stain for gangliosides.

Please replace the paragraph beginning at page 6, line

Applicants: Philip O. Livingston and Friedhelm Helling

Serial No.: 08/475,784 Filed: June 7, 1995

Page 3

24, with the following amended paragraph:

Figure 8B 8C

Inhibition of IgG reactivity of patient serum against GM2 and GD2. GM2 (A) and melanoma tissue ganglioside extract (B) were applied HPTLC plates, incubated with serum from patient No. 2 and stained with peroxidase-labeled goat antihuman IgG antibody. 3ml Patient serum at a dilution of 1:50 was preincubated with either  $150\mu g$  GM2 to  $150\mu g$  GD2 prior Immune staining.

Please replace the paragraph beginning at page 6, line 24, with the following amended paragraph:

Figures 9A, and 9B, 9C, and 9D

IqM and IgG antibody response in melanoma patients after immunization with GM2-KLH plus QS-21 vaccines. Sequential results for six patients receiving the 100 ug QS-21 dose are shown in Figures 9A and 9B Figure 9a and for six patients receiving the 200 ug dose in Figures 9C and 9D Figure 9b. Note that one patient in group received only vaccinations and was taken off study due to disease progression. Arrows indicate the time of cyclophosphamide (Cy) and GM2-KLH plus QS-21 vaccine injections.